

Appendix 2: Subgroup analyses in included studies (as supplied by the authors)

Table 1: Studies examining mortality among subgroups

Author	Subgroup	eGFR category	No. patients	HR (95% CI)	Adjustment variables
Traynor ¹	Nondiabetics	≥ 8 mL/min GFR < 8 mL/min GFR	97 87	Not reported	
Kazmi ²	Age > 67 y	Per ↑ 1 mL/min GFR	91 083	M1: 1.040 (1.038–1.042) M2: 1.028 (1.026–1.030) M3: 1.028 (1.025–1.031) M4: 1.028 (1.025–1.031)	M1: adjusted for age, sex, race, Hispanic ethnicity, BMI, cause of kidney failure, year of initiation of dialysis, network; M2: M1 + comorbid conditions; M3: M2 + hematocrit and albumin; M4: M3 + employment and insurance status
	Low-risk population	Per ↑ 1 mL/min GFR	90 540	M1: 1.047 (1.044–1.050) M2: 1.041 (1.038–1.044) M3: 1.034 (1.029–1.039) M4: 1.031 (1.026–1.036)	M1: adjusted for age, sex, race, Hispanic ethnicity, BMI, cause of kidney failure, year of initiation of dialysis, network; M2: M1 + comorbid conditions; M3: M2 + hematocrit and albumin; M4: M3 + employment and insurance status
Coronel ³	Diabetics on PD	> 7.7 mL/min/1.73 m ² ≤ 7.7 mL/min/1.73 m ²	56 44	Not reported	
Stel (1999 cohort) ⁴	Age 20–44 y 45–64 y 65–74 y > 75 y	Per ↑ 1 mL/min/1.73 m ²	4644†‡	1.04 (0.99–1.09) 1.05 (1.03–1.06) 1.04 (1.03–1.06) 1.03 (1.02–1.04)	Age at start of dialysis, gender, primary renal disease, treatment modality, and country
	Sex Female Male	Per ↑ 1 mL/min/1.73 m ²	4644	1.03 (1.01–1.05) 1.04 (1.03–1.05)	Same as above
	Comorbidity DM RVD/HTN GN Other	Per ↑ 1 mL/min/1.73 m ²	4644	1.04 (1.02–1.05) 1.05 (1.03–1.06) 1.03 (1.00–1.06) 1.05 (1.03–1.06)	Same as above
Stel (2003 cohort) ⁴	Age 20–44 y 45–64 y 65–74 y > 75 y	Per ↑ 1 mL/min/1.73 m ²	M1: 6613† M2: 3375‡	M1: 1.11 (1.05–1.17); M2: 1.12 (1.03–1.22) M1: 1.04 (1.02–1.06); M2: 1.05 (1.03–1.07) M1: 1.02 (1.01–1.04); M2: 1.02 (0.99–1.04) M1: 1.02 (1.01–1.03); M2: 1.01 (0.98–1.03)	M1: age at start of dialysis, gender, primary renal disease, treatment modality, and country; M2: M1 + diabetes, heart disease, PVD, cerebrovascular disease, malignancy
	Sex Female Male	Per ↑ 1 mL/min/1.73 m ²	M1: 6613 M2: 3375	M1: 1.02 (1.00–1.03); M2: 1.01 (0.99–1.04) M1: 1.03 (1.02–1.04); M2: 1.03 (1.01–1.05)	Same as above

Table 1: Studies examining mortality among subgroups

Author	Subgroup	eGFR category	No. patients	HR (95% CI)	Adjustment variables
	Comorbidity	Per ↑ 1 mL/min/1.73 m ²			Same as above
	DM		M1: 6613	M1: 1.02 (1.00–1.04); M2: 1.00 (0.99–1.04)	
	RVD/HTN		M2: 3375	M1: 1.02 (1.00–1.05); M2: 1.03 (1.00–1.07)	
	GN			M1: 1.10 (1.05–1.15); M2: 1.11 (1.05–1.18)	
	Other			M1: 1.02 (1.01–1.03); M2: 1.02 (1.00–1.04)	
	Dialysis	Per ↑ 1 mL/min/1.73 m ²			Same as above
	HD		M1: 6613	M1: 1.02 (1.02–1.04); M2: 1.02 (1.01–1.04)	
	PD		M2: 3375	M1: 1.03 (1.01–1.05); M2: 1.03 (1.00–1.05)	
Lassalle ⁵	Planned HD	Per ↑ 5 mL/min/1.73 m ²	6672	M1: 1.26 (1.19–1.34); M2: 1.13 (1.07–1.20); M3: 1.12 (1.06–1.19)	M1: age and gender; M2: M1 + diabetes, heart failure, dysrhythmia, PVD, CHD, malignancy, severe disability; M3: M2 + predialysis anemia care, initial treatment condition, wait listing or transplantation
	Planned PD		1367	M1: 1.16 (1.05–1.29); M2: 0.98 (0.88–1.10); M3: 1.00 (0.89–1.12)	
	Unplanned		3646	M1: 1.20 (1.13–1.26); 1.09 (1.03–1.15); M3: 1.08 (1.03–1.15)	
Cooper ⁶	Age				Unadjusted
	< 60 y	10–14	180	39/180§	
		5–7	194	38/194 (not significant)	
	≥ 60 y	10–14	224	113/224	
		5–7	230	117/230 (not significant)	
	Sex				
	Female	10–14	143	55/143	
		5–7	143	58/143 (not significant)	
	Male	10–14	261	97/261	
		5–7	281	97/281 (not significant)	
	Diabetes				
	No	10–14	232	65/232	
		5–7	241	63/241 (not significant)	
	Yes	10–14	172	87/172	
		5–7	183	92/183 (not significant)	
	Albumin				
	< 35 g/L	10–14	68	38/68	
		5–7	81	44/81 (not significant)	
	≥ 35 g/L	10–14	325	110/325	
		5–7	336	109/336 (not significant)	

Table 1: Studies examining mortality among subgroups

Author	Subgroup	eGFR category	No. patients	HR (95% CI)	Adjustment variables
Wright ⁷	Age	> 15	651 304	1.48 (1.46–1.49)	Age at ESRD onset, height and weight at ESRD onset, race, sex, diabetic status, Charlson comorbidity index, duration of predialysis nephrology care, type of dialysis, type of vascular access, cause of ESRD
		10–15		1.17 (1.16–1.17)	
		5–10		Reference	
		≤ 5		0.86 (0.85–0.86)	
		> 15	243 989	1.35 (1.33–1.37)	
		10–15		1.11 (1.10–1.26)	
		5–10		Reference	
		≤ 5		0.96 (0.94–0.97)	
	Dialysis	> 15	63 691	1.42 (1.37–1.47)	Same as above
		10–15		1.10 (1.07–1.12)	
		5–10		Reference	
		≤ 5		0.96 (0.93–0.99)	
		> 15	801 685	1.49 (1.47–1.50)	
		10–15		1.17 (1.16–1.17)	
		5–10		Reference	
		≤ 5		0.87 (0.86–0.87)	
Rosansky (2011) ⁸	Albumin ≥ 3.5 g/dL– HD only	< 6	204 208	1.46 (1.42–1.5)	Same as above
		10–15		1.18 (1.16–1.21)	
		5–10		Reference	
		≤ 5		0.84 (0.83–0.85)	
		6–8	468 446	1.46 (1.44–1.48)	Same as above
		10–15		1.15 (1.14–1.16)	
		5–10		Reference	
		≤ 5		0.90 (0.89–0.91)	
		> 8	222 639	1.37 (1.35–1.39)	Same as above
		10–15		1.13 (1.11–1.14)	
		5–10		Reference	
		≤ 5		0.94 (0.92–0.95)	

Table 1: Studies examining mortality among subgroups

Author	Subgroup	eGFR category	No. patients	HR (95% CI)	Adjustment variables
Kim ⁹	Dialysis	HD	< 5	47	Not detailed—(read off graph); significant difference at 60 months
			≥ 5	61	
	PD		< 5	52	Not detailed (read off graph); no significant difference at 60 months
			≥ 5	50	
Collins ¹⁰	HD	10–14	171	Reference	
		5–7	191	0.97 (0.66–1.41)	
Rosansky (2009) ¹¹	Age 65–74 y	5–9.9		Not provided	
		> 15			
Wilson ¹²	HD only	< 5.0	46	Reference	
		5.0–10.0	180	1.58 (0.54–4.65)¶	Sex, age, months at CKD clinic, comorbidity (cardiac disease, PVD, diabetes, antihypertensive use)
		> 10.0	45	1.68 (0.65–4.32)	
Tang ¹³	PD only	Per ↓ 1 mL/min/1.73 m ²	233	1.53 (1.20–3.99)	Age, gender, diabetic status
Shiao ¹⁴	PD only	Per ↑ 1 mL/min/1.73 m ²	275	1.18 (1.02–1.37)	Age, sex, level of education, occupational activity, nephrology referral, implantation of catheters, initiation of dialysis, comorbidities (diabetes, CAD, congestive heart failure), lab data
Hwang ¹⁵	HD only	Per ↑ 1 mL/min/1.73 m ²	23 551	1.15 (1.14–1.17)	Age, sex, diabetes, GN, HTN, chronic TIN, CAD, CHF CVD, malignancy, liver cirrhosis, TB, dialysis initiation year
Clark ¹⁶	HD only	Per ↑ 1 mL/min/1.73 m ²	25 910	1.01 (1.01–1.02)	Age, sex, ethnicity, DM, GN, RVD, modified CMI score, CAD, CHR, HTN, CVD, PVD, lung disease, malignancy, albumin, vascular access, late referral

Note: BMI = body mass index; CAD = coronary artery disease; CHD: coronary heart disease; CHF = congestive heart failure; CKD = chronic kidney disease; CMi= Charlson Comorbidity Index; CVD = cardiovascular disease; DM = diabetes mellitus; eGFR = estimated glomerular filtration rate; ESRD = end-stage renal disease; GN = glomerulonephritis; HD = hemodialysis; HR = hazard ratio; HTN = hypertension; M = model; PD = peritoneal dialysis; PVD = peripheral vascular disease; RVD = renal vascular disease; TB = tuberculosis; TIN = tubulointerstitial nephritis.

*Without diabetes, congestive heart failure, or heart disease.

†Not presenting unadjusted analyses.

#Number of patients in each subgroup not detailed for each model.

§Hazard ratio not written out (read off of table).

||Confidence intervals not presented.

¶Odds ratio, year 2 mortality (year 1 mortality not presented).

References

1. Traynor J, Simpson K, Geddes C, Deighan C, Fox J. Early initiation of dialysis fails to prolong survival in patients with end-stage renal failure. *J Am Soc Nephrol*. 2002;13:2125-32.
2. Kazmi W, Gilbertson D, Obrador G, et al. Effect of comorbidity on the increased mortality associated with early initiation of dialysis. *Am J Kidney Dis* 2005;46:887-96.
3. Coronel F, Cigarran S, Herrero J. Early initiation of peritoneal dialysis in diabetic patients. *Scand J Urol Nephrol* 2009;43:148-53.
4. Stel V, Dekker F, Ansell D, et al. Residual renal function at the start of dialysis and clinical outcomes. *Nephrol Dial Transplant* 2009;24:3175-82.
5. Lassalle M, Labeeuw M, Frimat L, et al. Age and comorbidity may explain the paradoxical association of an early dialysis start with poor survival. *Kidney Int* 2010;77.
6. Cooper B, Branley P, Bulfone L, Collins JF, Craig JC, Fraenkel MB, et al. A randomized, controlled trial of early versus late initiation of dialysis. *N Engl J Med* 2010;363:609-19.
7. Wright S, Klausner D, Baird B, et al. Timing of dialysis initiation and survival in ESRD. *Clin J Am Soc Nephrol* 2010;5:1828-35.
8. Rosansky S, Eggers P, Jackson K, Glasscock R, Clark W. Early start of hemodialysis may be harmful. *Arch Intern Med* 2011;17:396-403.
9. Kim S, Kim N. The effect of residual renal function at the initiation of dialysis on patient survival. *Korean J Intern Med* 2009;24:55-62.
10. Collins J, Cooper B, Branley P, et al. Outcomes of patients with planned initiation of hemodialysis in the IDEAL trial. *Contrib Nephrol* 2011;171:1-9.
11. Rosansky S, Clark W, Eggers P, Glasscock R. Initiation of dialysis at higher GFRs: is the apparent rising tide of early dialysis harmful or helpful? *Kidney Int* 2009;76:257-61.
12. Wilson B, Harwood L, Locking-Cusolito H, et al. Optimal timing of initiation of chronic hemodialysis? *Hemodial* 2007;11:263-9.
13. Tang S, Ho Y, Tang A, et al. Delaying initiation of dialysis till symptomatic uraemia—is it too late? *Nephrol Dial Transplant* 2007;22:1926-32.
14. Shiao C, Huang J, Chien K, Chuang H, Chen Y, Wu K. Early initiation of dialysis and late implantation of catheters adversely affect outcomes of patients on chronic peritoneal dialysis. *Perit Dial Int* 2008;28:73-81.
15. Hwang S-J, Yang W-C, Lin M-Y, May L-W, Chen H-C, Taiwan Society of Nephrology. Impact of the clinical conditions at dialysis initiation on mortality in incident haemodialysis patients: a national cohort study in Taiwan. *Nephrol Dial Transplant* 2010;25:2616-24.
16. Clark W, Na Y, Rosansky S, et al. Association between estimated glomerular filtration rate at initiation of dialysis and mortality. *CMAJ* 2011;183:47-54.